

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Juluca 50 mg/25 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains dolutegravir sodium equivalent to 50 mg dolutegravir and rilpivirine hydrochloride equivalent to 25 mg rilpivirine.

### Excipient with known effect

Each film-coated tablet contains 52 mg lactose (as monohydrate).

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Pink, oval, biconvex tablets, approximately 14 x 7 mm, debossed with 'SV J3T' on one side.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Juluca is indicated for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adults who are virologically-suppressed (HIV-1 RNA <50 copies/mL) on a stable antiretroviral regimen for at least six months with no history of virological failure and no known or suspected resistance to any non-nucleoside reverse transcriptase inhibitor or integrase inhibitor (see section 5.1).

### 4.2 Posology and method of administration

Juluca should be prescribed by physicians experienced in the management of HIV infection.

#### Posology

The recommended dose of Juluca is one tablet once daily. Juluca must be taken with a meal (see section 5.2).

Separate preparations of dolutegravir or rilpivirine are available in cases where discontinuation or dose adjustment of one of the active substances is indicated (see section 4.5). In these cases the physician should refer to the Summary of Product Characteristics for these medicinal products.

#### *Missed doses*

If the patient misses a dose of Juluca, the patient should take Juluca with a meal as soon as possible, providing the next dose is not due within 12 hours. If the next dose is due within 12 hours, the patient should not take the missed dose and simply resume the usual dosing schedule.

If a patient vomits within 4 hours of taking Juluca, another Juluca tablet should be taken with a meal. If a patient vomits more than 4 hours after taking Juluca, the patient does not need to take another dose of Juluca until the next regularly scheduled dose.

#### *Elderly*

There are limited data available on the use of Juluca in patients aged 65 years and over. There is no evidence that elderly patients require a different dose than younger adult patients (see section 5.2).

#### *Renal impairment*

No dosage adjustment is required in patients with mild or moderate renal impairment. In patients with severe renal impairment or end stage renal disease, the combination of Juluca with a strong CYP3A inhibitor should only be used if the benefit outweighs the risk. No data are available in subjects receiving dialysis although differences in pharmacokinetics are not expected in this population (see section 5.2).

#### *Hepatic impairment*

No dosage adjustment is required in patients with mild or moderate hepatic impairment (Child-Pugh score A or B). Juluca should be used with caution in patients with moderate hepatic impairment. No data are available in patients with severe hepatic impairment (Child-Pugh score C); therefore Juluca is not recommended in these patients (see section 5.2).

#### *Paediatric population*

The safety and efficacy of Juluca in children and adolescents aged less than 18 years have not yet been established. Currently available data are described in section 5.2, but no recommendation on a posology can be made.

#### *Pregnancy*

The safety and efficacy of Juluca in pregnancy have not yet been established. Limited data are available regarding the use of dolutegravir during pregnancy. Lower exposures of dolutegravir and rilpivirine were observed during pregnancy. No recommendations for dose adjustments can be made for Juluca. Therefore, use of Juluca during pregnancy is not recommended (see sections 4.4, 4.6, 5.1 and 5.2).

### Method of administration

#### *Oral use*

Juluca must be taken orally, once daily **with a meal** (see section 5.2). It is recommended that the film-coated tablet be swallowed whole with water and not be chewed or crushed.

## **4.3 Contraindications**

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

Co-administration with the following medicinal products:

- fampridine (also known as dalfampridine);
- carbamazepine, oxcarbazepine, phenobarbital, phenytoin;
- rifampicin, rifapentine;
- proton pump inhibitors, such as omeprazole, esomeprazole, lansoprazole, pantoprazole, rabeprazole;
- systemic dexamethasone, except as a single dose treatment;
- St John's wort (*Hypericum perforatum*).

## **4.4 Special warnings and precautions for use**

### Transmission of HIV

While effective viral suppression with antiretroviral therapy has been proven to substantially reduce the risk of sexual transmission, a residual risk cannot be excluded. Precautions to prevent transmission should be taken in accordance with national guidelines.

#### Hypersensitivity reactions

Hypersensitivity reactions have been reported with dolutegravir, and were characterized by rash, constitutional findings, and sometimes, organ dysfunction, including severe liver reactions. Juluca should be discontinued immediately if signs or symptoms of hypersensitivity reactions develop (including, but not limited to, severe rash or rash accompanied by raised liver enzymes, fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, facial oedema, eosinophilia, angioedema). Clinical status including liver aminotransferases and bilirubin should be monitored. Delay in stopping treatment with Juluca after the onset of hypersensitivity may result in a life-threatening allergic reaction.

#### Cardiovascular

At supra-therapeutic doses (75 and 300 mg once daily), rilpivirine has been associated with prolongation of the QTc interval of the electrocardiogram (ECG) (see sections 4.5 and 5.1). Rilpivirine at the recommended dose of 25 mg once daily is not associated with a clinically relevant effect on QTc. Juluca should be used with caution when co-administered with medicinal products with a known risk of Torsade de Pointes.

#### Opportunistic infections

Patients should be advised that Juluca does not cure HIV infection and that they may still develop opportunistic infections and other complications of HIV infection. Therefore, patients should remain under close clinical observation by physicians experienced in the treatment of these associated HIV diseases.

#### Osteonecrosis

Although the aetiology is considered to be multifactorial (including corticosteroid use, biphosphonates, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported in patients with advanced HIV-disease and/or long-term exposure to CART. Patients should be advised to seek medical advice if they experience joint aches and pain, joint stiffness or difficulty in movement.

#### Patients with hepatitis B or C

No clinical data are available in patients with hepatitis B co-infection. Physicians should refer to current treatment guidelines for the management of HIV infection in patients co-infected with hepatitis B virus. Limited data is available in patients with hepatitis C co-infection. A higher incidence of liver chemistry elevations (Grade 1) were observed in patients treated with dolutegravir and rilpivirine co-infected with hepatitis C compared to those who were not co-infected. Monitoring of liver function is recommended in patients with hepatitis B and/or C co-infection.

#### Interactions with other medicinal products

Juluca should not be administered with other antiretroviral medicinal products for the treatment of HIV (see section 4.5).

Juluca should not be co-administered at the same time as H<sub>2</sub>-receptor antagonists. These medicinal products are recommended to be administered 12 hours before or 4 hours after Juluca (see section 4.5).

Juluca should not be co-administered at the same time as antacids. These medicinal products are recommended to be administered 6 hours before or 4 hours after Juluca (see section 4.5).

Calcium or iron supplements, or multivitamins should be co-administered at the same time as Juluca, with a meal. If calcium or iron supplements, or multivitamins cannot be taken at the same time as Juluca, these supplements are recommended to be administered 6 hours before or 4 hours after taking Juluca (see section 4.5).

Dolutegravir increased metformin concentrations. A dose adjustment of metformin should be considered when starting and stopping co-administration of Juluca with metformin, to maintain glycaemic control (see section 4.5). Metformin is eliminated renally and therefore it is of importance to monitor renal function when co-treated with Juluca. This combination may increase the risk for lactic acidosis in patients with moderate renal impairment (stage 3a creatinine clearance [CrCl] 45– 59 mL/min) and a cautious approach is recommended. Reduction of the metformin dose should be highly considered.

Juluca should not be taken with any other medicinal product containing dolutegravir or rilpivirine, except in case of co-administration with rifabutin (see section 4.5).

### Pregnancy

The safety and efficacy of Juluca in pregnancy have not yet been established. Limited data are available regarding the use of dolutegravir during pregnancy. Lower exposures of dolutegravir or rilpivirine were observed when taken once daily, in combination with a background regimen, during pregnancy. In phase 3 studies, lower rilpivirine exposure, similar to that seen during pregnancy, has been associated with an increased risk of virological failure. No recommendations for dose adjustments can be made for Juluca. Therefore, use of Juluca during pregnancy is not recommended (see sections 4.6, 5.1 and 5.2).

### Immune Reactivation Syndrome

In HIV-infected patients with severe immune deficiency at the time of institution of combination antiretroviral therapy (CART), an inflammatory reaction to asymptomatic or residual opportunistic pathogens may arise and cause serious clinical conditions, or aggravation of symptoms. Typically, such reactions have been observed within the first few weeks or months of initiation of CART. Relevant examples are cytomegalovirus retinitis, generalised and/or focal mycobacterial infections, and Pneumocystis jirovecii pneumonia. Any inflammatory symptoms should be evaluated and treatment instituted when necessary. Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution, however, the reported time to onset is more variable and these events can occur many months after initiation of treatment.

### Excipients

Juluca contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Juluca is intended for use as a complete regimen for the treatment of HIV-1 infection and should not be administered with other antiretroviral medicinal products for the treatment of HIV. Therefore, information regarding drug-drug interactions with other antiretroviral medicinal products is not provided. Juluca contains dolutegravir and rilpivirine, therefore any interactions identified with these active substances are relevant to Juluca. Interaction studies have only been performed in adults.

### Effect of other medicinal products on the pharmacokinetics of dolutegravir and rilpivirine

Dolutegravir is eliminated mainly through metabolism by uridine diphosphate glucuronosyl transferase (UGT)1A1. Dolutegravir is also a substrate of UGT1A3, UGT1A9, cytochrome P450 (CYP)3A4, P-glycoprotein (P-gp), and breast cancer resistance protein (BCRP); therefore medicinal products that induce those enzymes may decrease dolutegravir plasma concentration and reduce the therapeutic effect of

dolutegravir (see Table 1). Co-administration of Juluca and other medicinal products that inhibit these enzymes may increase dolutegravir plasma concentration (see Table 1).

The absorption of dolutegravir is reduced by certain anti-acid medicinal products (see Table 1).

Rilpivirine is primarily metabolised by CYP3A. Medicinal products that induce or inhibit CYP3A may thus affect the clearance of rilpivirine (see section 5.2). Co-administration of Juluca with medicinal products that induce CYP3A may result in decreased plasma concentrations of rilpivirine, which could reduce the therapeutic effect of Juluca (see Table 1). Co-administration of Juluca with medicinal products that inhibit CYP3A may result in increased plasma concentrations of rilpivirine (see Table 1).

Co-administration of Juluca with medicinal products that increase gastric pH may result in decreased plasma concentrations of rilpivirine which could potentially reduce the therapeutic effect of Juluca.

#### Effect of dolutegravir and rilpivirine on the pharmacokinetics of other medicinal products

Based on *in vivo* and/or *in vitro* data, dolutegravir is not expected to affect the pharmacokinetics of medicinal products that are substrates of any major enzyme or transporter such as CYP3A4, CYP2C9 and P-gp (for more information see section 5.2).

*In vitro*, dolutegravir inhibited the renal organic cation transporter 2 (OCT2) and multidrug and toxin extrusion transporter 1 (MATE1). *In vivo*, a 10-14% decrease of creatinine clearance (secretory fraction is dependent on OCT2 and MATE1 transport) was observed in patients. *In vivo*, dolutegravir may increase plasma concentrations of medicinal products in which excretion is dependent upon OCT2 and/or MATE1 (e.g. fampridine [also known as dalfampridine], metformin) (see Table 1 and sections 4.3 and 4.4).

*In vitro*, dolutegravir inhibited the renal uptake transporters, organic anion transporters (OAT)1 and OAT3. Based on the lack of effect on the *in vivo* pharmacokinetics of the OAT substrate tenofovir, *in vivo* inhibition of OAT1 is unlikely. Inhibition of OAT3 has not been studied *in vivo*. Dolutegravir may increase plasma concentrations of medicinal products in which excretion is dependent upon OAT3.

Rilpivirine 25 mg once daily is not likely to have a clinically relevant effect on the exposure of medicinal products metabolised by CYP enzymes.

Rilpivirine inhibits P-gp *in vitro* ( $IC_{50}$  is 9.2  $\mu\text{M}$ ). In a clinical study, rilpivirine did not significantly affect the pharmacokinetics of digoxin. However, it may not be completely excluded that rilpivirine can increase the exposure to other medicinal products transported by P-gp that are more sensitive to intestinal P-gp inhibition, e.g. dabigatran etexilate.

Rilpivirine is an *in vitro* inhibitor of the transporter MATE-2K with an  $IC_{50}$  of < 2.7 nM. The clinical implications of this finding are currently unknown.

#### Interaction table

Selected established and theoretical interactions between dolutegravir, rilpivirine and co-administered medicinal products are listed in Table 1.

(increase is indicated as “↑”, decrease as “↓”, no change as “↔”, area under the concentration versus time curve as “AUC”, maximum observed concentration as “ $C_{\max}$ ”, minimum observed concentration as “ $C_{\min}$ ” concentration at end of dosing interval as “ $C_t$ ”).

**Table 1:** **Drug Interactions**

Medicinal products by therapeutic areas	Interaction Geometric mean change (%)	Recommendations concerning co-administration
---	---------------------------------------	--

<b>Antiviral active substances</b>		
Tenofovir disoproxil / Dolutegravir <sup>1</sup>	Dolutegravir ↔ AUC ↑ 1% $C_{\max} \downarrow 3\%$ $C\tau \downarrow 8\%$  Tenofovir ↔	No dose adjustment is required.
Tenofovir disoproxil / Rilpivirine <sup>1,2</sup>	Rilpivirine AUC ↔ $C_{\min} \leftrightarrow$ $C_{\max} \leftrightarrow$  Tenofovir AUC ↑ 23% $C_{\min} \uparrow 24\%$ $C_{\max} \uparrow 19\%$	
Tenofovir alafenamide / Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Tenofovir alafenamide / Rilpivirine <sup>1</sup>	Rilpivirine ↔	
Lamivudine/ Dolutegravir	Dolutegravir ↔	No dose adjustment is required.
Lamivudine/ Rilpivirine	Rilpivirine ↔ (Not studied)	
Entecavir/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Entecavir/ Rilpivirine	Rilpivirine ↔ (Not studied)	
Daclatasvir/ Dolutegravir <sup>1</sup>	Dolutegravir ↔ AUC ↑ 33% $C_{\max} \uparrow 29\%$ $C\tau \uparrow 45\%$  Daclatasvir ↔	No dose adjustment is required.
Daclatasvir/ Rilpivirine	Rilpivirine ↔	
Simeprevir/ Dolutegravir	Dolutegravir ↔	No dose adjustment is required.
Simeprevir/ Rilpivirine	Rilpivirine ↔ AUC ↔ $C_{\min} \uparrow 25\%$ $C_{\max} \leftrightarrow$ Simeprevir ↔ AUC ↔ $C_{\min} \leftrightarrow$ $C_{\max} \uparrow 10\%$	

Sofosbuvir / Dolutegravir <sup>1</sup>	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Sofosbuvir / Rilpivirine	Rilpivirine ↔ AUC ↔ $C_{\min}$ ↔ $C_{\max}$ ↔ Sofosbuvir ↔ AUC ↔ $C_{\max} \uparrow 21\%$ Sofosbuvir metabolite GS- 331007 ↔ AUC ↔ $C_{\max}$ ↔	
Ledipasvir/Sofosbuvir / Dolutegravir <sup>1</sup>	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Ledipasvir/Sofosbuvir / Rilpivirine	Rilpivirine ↔ AUC ↓ 5% $C_{\min} \downarrow 7\%$ $C_{\max} \downarrow 3\%$ Ledipasvir ↔ AUC ↑ 2% $C_{\min} \uparrow 2\%$ $C_{\max} \uparrow 1\%$ Sofosbuvir ↔ AUC ↑ 5% $C_{\max} \downarrow 4\%$ Sofosbuvir metabolite GS- 331007 ↔ AUC ↑ 8% $C_{\min} \uparrow 10\%$ $C_{\max} \uparrow 8\%$	
Sofosbuvir/ Velpatasvir/ Dolutegravir <sup>1</sup>	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Sofosbuvir/ Velpatasvir/ Rilpivirine	Rilpivirine ↔ AUC ↔ $C_{\min}$ ↔ $C_{\max}$ ↔ Sofosbuvir ↔ AUC ↔ $C_{\max}$ ↔ Sofosbuvir metabolite GS- 331007 ↔ AUC ↔ $C_{\min}$ ↔ $C_{\max}$ ↔ Velpatasvir ↔ AUC ↔ $C_{\min}$ ↔ $C_{\max}$ ↔	
Ribavirin/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.

Ribavirin/ Rilpivirine	Rilpivirine ↔ (Not studied)	
<b>Other active substances</b>		
<i>Antiarrhythmics</i>		
Digoxin/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Digoxin/ Rilpivirine <sup>1</sup>	Rilpivirine ↔  Digoxin AUC ↔ $C_{min}$ NA $C_{max}$ ↔	
<i>Anticonvulsants</i>		
Carbamazepine/ Dolutegravir <sup>1</sup>	Dolutegravir ↓ AUC ↓ 49% $C_{max}$ ↓ 33% $C\tau$ ↓ 73%	Metabolic inducers may significantly decrease dolutegravir/rilpivirine plasma concentrations, resulting in loss of therapeutic effect. Co-administration of Juluca with these metabolic inducers is contraindicated (see section 4.3).
Carbamazepine/ Rilpivirine	Rilpivirine ↓ Not studied. Significant decreases in rilpivirine plasma concentrations are expected (induction of CYP3A enzymes).	
Oxcarbazepine Phenytoin Phenobarbital/ Dolutegravir	Dolutegravir ↓ Not studied. Decrease expected due to induction of UGT1A1 and CYP3A enzymes, a similar reduction in exposure as observed with carbamazepine is expected.	Metabolic inducers may significantly decrease dolutegravir/rilpivirine plasma concentrations, resulting in loss of therapeutic effect. Co-administration of Juluca with these metabolic inducers is contraindicated (see section 4.3).
Oxcarbazepine Phenytoin Phenobarbital/ Rilpivirine	Rilpivirine ↓ Not studied. Significant decreases in rilpivirine plasma concentrations are expected (induction of CYP3A enzymes).	
<i>Azole anti-fungals</i>		
Ketoconazole/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Ketoconazole/ Rilpivirine <sup>1,2</sup>	Rilpivirine AUC ↑ 49% $C_{min}$ ↑ 76% $C_{max}$ ↑ 30% (inhibition of CYP3A enzymes).  Ketoconazole AUC ↓ 24%	

	$C_{min} \downarrow 66\%$ $C_{max} \leftrightarrow$ (induction of CYP3A due to high rilpivirine dose in the study).	
Fluconazole Itraconazole Isavuconazole Posaconazole Voriconazole/ Dolutegravir	Dolutegravir $\leftrightarrow$ (Not studied)	No dose adjustment is required.
Fluconazole Itraconazole Isavuconazole Posaconazole Voriconazole/ Rilpivirine	Rilpivirine $\uparrow$ Not studied. May cause an increase in the plasma concentrations of rilpivirine (inhibition of CYP3A enzymes).	
<i>Herbal products</i>		
St. John's wort/ Dolutegravir	Dolutegravir $\downarrow$ Not studied. Decrease expected due to induction of UGT1A1 and CYP3A enzymes, a similar reduction in exposure as observed with carbamazepine is expected.	Co-administration may cause significant decreases in rilpivirine plasma concentrations. This may result in loss of therapeutic effect of Juluca. Co-administration of Juluca with St. John's wort is contraindicated (see section 4.3).
St. John's wort/ Rilpivirine	Rilpivirine $\downarrow$ Not studied. Significant decreases in rilpivirine plasma concentrations are expected (induction of CYP3A enzymes).	
<i>Potassium channel blockers</i>		
Fampridine (also known as dalfampridine)/ Dolutegravir	Fampridine $\uparrow$	Co-administration of dolutegravir has the potential to cause seizures due to increased fampridine plasma concentration via inhibition of OCT2 transporter; co-administration has not been studied. Fampridine co-administration with Juluca is contraindicated (see section 4.3).
<i>Proton pump inhibitors</i>		
Omeprazole Lansoprazole Rabeprazole Pantoprazole Esomeprazole/ Dolutegravir	Dolutegravir $\leftrightarrow$ (Not studied)	Co-administration may significantly decrease rilpivirine plasma concentration. This may result in loss of therapeutic effect of Juluca. Co-administration of Juluca with proton pump inhibitors is contraindicated (see section 4.3).
Omeprazole/ Rilpivirine <sup>1,2</sup>	Rilpivirine $AUC \downarrow 40\%$ $C_{min} \downarrow 33\%$	

	$C_{max} \downarrow 40\%$ (reduced absorption due to gastric pH increase).	
	Omeprazole AUC $\downarrow 14\%$ $C_{min}$ NA $C_{max} \downarrow 14\%$	
Lansoprazole Rabeprazole Pantoprazole Esomeprazole/ Rilpivirine	Rilpivirine $\downarrow$ Not studied. Significant decreases in rilpivirine plasma concentrations are expected (reduced absorption due to gastric pH increase).	

#### *H<sub>2</sub>-receptor antagonists*

Famotidine Cimetidine Nizatidine Ranitidine/ Dolutegravir	Dolutegravir $\leftrightarrow$ (Not studied)	The combination of Juluca and H <sub>2</sub> -receptor antagonists should be used with particular caution. Only H <sub>2</sub> -receptor antagonists that can be dosed once daily should be used.
Famotidine/ Rilpivirine <sup>1,2</sup> 40 mg single dose taken 12 hours before rilpivirine	Rilpivirine AUC $\downarrow 9\%$ $C_{min}$ NA $C_{max} \leftrightarrow$	H <sub>2</sub> -receptor antagonists should be taken well separated in time from the administration of Juluca (minimum 4 hours after or 12 hours before)
Famotidine/ Rilpivirine <sup>1,2</sup> 40 mg single dose taken 2 hours before rilpivirine	Rilpivirine AUC $\downarrow 76\%$ $C_{min}$ NA $C_{max} \downarrow 85\%$ (reduced absorption due to gastric pH increase).	
Famotidine/ Rilpivirine <sup>1,2</sup> 40 mg single dose taken 4 hours after rilpivirine	Rilpivirine AUC $\uparrow 13\%$ $C_{min}$ NA $C_{max} \uparrow 21\%$	

#### *Antacids and supplements*

Antacids (e.g., aluminium magnesium hydroxide, and/or calcium)	Dolutegravir $\downarrow$ AUC $\downarrow 74\%$ $C_{max} \downarrow 72\%$ $C_{24} \downarrow 74\%$	The combination of Juluca and antacids should be used with particular caution. Antacids should be taken well separated in time from the
--	---	---

carbonate)/ Dolutegravir <sup>1</sup>	(Complex binding to polyvalent ions).	administration of Juluca (minimum 6 hours before or 4 hours after).
Antacids (e.g., aluminium magnesium hydroxide, and/or calcium carbonate)/ Rilpivirine	Rilpivirine ↓ Not studied. Significant decreases in rilpivirine plasma concentrations are expected (reduced absorption due to gastric pH increase).	
Calcium supplements/ Dolutegravir <sup>1</sup>	Dolutegravir ↓ AUC ↓ 39% $C_{max}$ ↓ 37% $C_{24}$ ↓ 39% (Complex binding to polyvalent ions).	The combination of Juluca and supplements should be used with particular caution. Calcium supplements, iron supplements or multivitamins should be co-administered at the same time as Juluca with a meal.
Iron supplements/ Dolutegravir <sup>1</sup>	Dolutegravir ↓ AUC ↓ 54% $C_{max}$ ↓ 57% $C_{24}$ ↓ 56% (Complex binding to polyvalent ions).	If calcium supplements, iron supplements or multivitamins cannot be taken at the same time as Juluca, these supplements should be taken well separated in time from the administration of Juluca (minimum 6 hours before or 4 hours after).
Multivitamin/ Dolutegravir <sup>1</sup>	Dolutegravir ↓ AUC ↓ 33% $C_{max}$ ↓ 35% $C_{24}$ ↓ 32% (Complex binding to polyvalent ions).	
<b>Corticosteroids</b>		
Prednisone/ Dolutegravir <sup>1</sup>	Dolutegravir ↔ AUC ↑ 11% $C_{max}$ ↑ 6% $C\tau$ ↑ 17%	No dose adjustment is required.
Prednisone/ Rilpivirine	Rilpivirine ↔ (Not studied)	
Dexamethasone/ Dolutegravir	Dolutegravir ↔ (Not studied)	Co-administration may cause significant decreases in rilpivirine plasma concentrations. This may result in loss of therapeutic effect of Juluca.
Dexamethasone/ Rilpivirine (systemic, except for single dose use)	Rilpivirine ↓ Not studied. Dose dependent decreases in rilpivirine plasma concentrations are expected (induction of CYP3A enzymes).	Co-administration of Juluca with systemic dexamethasone is contraindicated (except as a single dose) see section 4.3. Alternatives should be considered, particularly for long-term use.
<b>Antidiabetics</b>		
Metformin/ Dolutegravir <sup>1</sup>	Metformin ↑ AUC ↑ 79% $C_{min}$ NA $C_{max}$ ↑ 66%	A dose adjustment of metformin should be considered when starting and stopping co-administration of Juluca with metformin, to maintain glycaemic control. In patients with moderate renal impairment a dose adjustment of metformin should be considered when co-administered with dolutegravir, because of the increased risk for lactic acidosis in patients with
Metformin/ Rilpivirine <sup>1</sup>	Metformin AUC ↔ $C_{min}$ NA	

	$C_{\max} \leftrightarrow$	moderate renal impairment due to increased metformin concentration (section 4.4).
<i>Antimycobacterials</i>		
Rifampicin/ Dolutegravir <sup>1</sup>	Dolutegravir ↓ AUC ↓ 54% $C_{\max} \downarrow 43\%$ $C_{\tau} \downarrow 72\%$ (induction of UGT1A1 and CYP3A enzymes).	Co-administration may cause significant decreases in rilpivirine plasma concentrations. This may result in loss of therapeutic effect of Juluca. Co-administration of Juluca with rifampicin is contraindicated (see section 4.3).
Rifampicin/ Rilpivirine <sup>1,2</sup>	Rilpivirine AUC ↓ 80% $C_{\min} \downarrow 89\%$ $C_{\max} \downarrow 69\%$ (induction of CYP3A enzymes).  Rifampicin AUC ↔ $C_{\min}$ NA $C_{\max} \leftrightarrow$ 25-desacetyl-rifampicin AUC ↓ 9% $C_{\min}$ NA $C_{\max} \leftrightarrow$	
Rifabutin/ Dolutegravir <sup>1</sup>	Dolutegravir ↔ AUC ↓ 5% $C_{\max} \uparrow 16\%$ $C_{\tau} \downarrow 30\%$ (induction of UGT1A1 and CYP3A enzymes).	Co-administration is likely to cause significant decreases in rilpivirine plasma concentrations (induction of CYP3A enzymes). When Juluca is co-administered with rifabutin, an additional 25 mg tablet of rilpivirine per day should be taken at the same time with Juluca, for the duration of the rifabutin co-administration (a separate formulation of rilpivirine is available for this dose adjustment, see section 4.2).
Rifabutin/ Rilpivirine <sup>1</sup> 300 mg once daily <sup>2</sup>	Rifabutin AUC ↔ $C_{\min} \leftrightarrow$ $C_{\max} \leftrightarrow$ 25-O-desacetyl-rifabutin AUC ↔ $C_{\min} \leftrightarrow$ $C_{\max} \leftrightarrow$	
300 mg once daily (+ 25 mg once daily rilpivirine)	Rilpivirine AUC ↓ 42% $C_{\min} \downarrow 48\%$ $C_{\max} \downarrow 31\%$	
300 mg once daily (+ 50 mg once daily rilpivirine)	Rilpivirine AUC ↑ 16%* $C_{\min} \leftrightarrow^*$ $C_{\max} \uparrow 43\%*$	
	* compared to 25 mg once daily rilpivirine alone	

	(induction of CYP3A enzymes).	
Rifapentine/ Dolutegravir	Dolutegravir ↓ (Not studied)	Co-administration may cause significant decreases in rilpivirine plasma concentrations. This may result in loss of therapeutic effect of Juluca (induction of CYP3A enzymes).
Rifapentine/ Rilpivirine	Rilpivirine ↓ Not studied. Significant decreases in rilpivirine plasma concentrations are expected.	Co-administration of Juluca with rifapentine is contraindicated (see section 4.3).
<i>Antimalarials</i>		
Artemether/ Lumefantrine/ Dolutegravir	Dolutegravir ↔ (Not studied)	The combination of Juluca and artemether/lumefantrine should be used with caution.
Artemether/ Lumefantrine/ Rilpivirine	Rilpivirine ↓ Not studied. Decreased exposure of rilpivirine is expected (induction of CYP3A enzymes).	
Atovaquone/ Proguanil/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Atovaquone/ Proguanil/ Rilpivirine	Rilpivirine ↔ (Not studied).	
<i>Macrolide antibiotics</i>		
Clarithromycin Erythromycin /Dolutegravir	Dolutegravir ↔ (Not studied)	Where possible, alternatives such as azithromycin should be considered.
Clarithromycin Erythromycin /Rilpivirine	Rilpivirine ↑ Not studied. Increased exposure of rilpivirine is expected (inhibition of CYP3A enzymes).	
<i>Oral contraceptives</i>		
Ethinyl estradiol (EE) <sup>1</sup> and Norelgestromin (NGMN) <sup>1</sup> / Dolutegravir	Dolutegravir ↔ EE ↔ AUC ↑ 3% $C_{max}$ ↓ 1%  NGMN ↔ AUC ↓ 2% $C_{max}$ ↓ 11%	Dolutegravir or rilpivirine did not change ethinyl estradiol and norelgestromin (dolutegravir) or norethindrone (rilpivirine) plasma concentrations to a clinically relevant extent. No dose adjustment of oral contraceptives is required when co-administered with Juluca.
Ethinyl estradiol (EE) <sup>1</sup> and Norethindrone <sup>1</sup> / Rilpivirine	Rilpivirine ↔* EE ↔ AUC ↔ $C_{min}$ ↔ $C_{max}$ ↑ 17%	

	Norethindrone ↔ AUC ↔ $C_{\min}$ ↔ $C_{\max}$ ↔  *based on historic controls.	
<i>Analgesics</i>		
Methadone/ Dolutegravir <sup>1</sup>	Dolutegravir ↔ Methadone ↔ AUC ↓ 2% $C_{\max}$ ↔ 0% $C_{\tau}$ ↓ 1%	No dose adjustments are required when initiating co-administration of methadone with Juluca. However, clinical monitoring is recommended as methadone maintenance therapy may need to be adjusted in some patients.
Methadone / Rilpivirine <sup>1</sup>	Rilpivirine: AUC: ↔* $C_{\min}$ : ↔* $C_{\max}$ : ↔*  R(-) methadone: AUC: ↓ 16% $C_{\min}$ : ↓ 22% $C_{\max}$ : ↓ 14%  *based on historic controls.	
Paracetamol/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Paracetamol / Rilpivirine <sup>1,2</sup>	Rilpivirine AUC ↔ $C_{\min}$ ↑ 26% $C_{\max}$ ↔  Paracetamol AUC ↔ $C_{\min}$ NA $C_{\max}$ ↔	
<i>Anticoagulants</i>		
Dabigatran etexilate/ Dolutegravir	Dolutegravir ↔ (Not studied)	The combination of Juluca and dabigatran etexilate should be used with caution.
Dabigatran etexilate/ Rilpivirine	Rilpivirine ↔ Not studied. Dabigatran etexilate ↑ A risk for increases in dabigatran plasma concentrations cannot be excluded (inhibition of intestinal P-gp).	
<i>HMG CO-A reductase inhibitors</i>		
Atorvastatin/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
	Rilpivirine	

Atorvastatin/ Rilpivirine <sup>1,2</sup>	AUC ↔ $C_{min} \leftrightarrow$ $C_{max} \downarrow 9\%$  Atorvastatin AUC ↔ $C_{min} \downarrow 15\%$ $C_{max} \uparrow 35\%$	
<i>Phosphodiesterase type 5 (PDE-5) inhibitors</i>		
Sildenafil / Dolutegravir	Dolutegravir ↔	No dose adjustment is required.
Sildenafil/ Rilpivirine <sup>1,2</sup>	Rilpivirine AUC ↔ $C_{min} \leftrightarrow$ $C_{max} \leftrightarrow$  Sildenafil AUC ↔ $C_{min} NA$ $C_{max} \leftrightarrow$	
Vardenafil Tadalafil/ Dolutegravir	Dolutegravir ↔ (Not studied)	No dose adjustment is required.
Vardenafil Tadalafil/ Rilpivirine	Rilpivirine ↔ (Not studied)	

<sup>1</sup> The interaction between dolutegravir and/or rilpivirine and the medicinal product was evaluated in a clinical study. All other drug-drug interactions shown are predicted.

<sup>2</sup> This interaction study has been performed with a dose higher than the recommended dose for rilpivirine assessing the maximal effect on the co-administered drug.

NA = Not applicable

#### QT prolonging medicinal products

There is limited information available on the potential for a pharmacodynamic interaction between rilpivirine and medicinal products that prolong the QTc interval of the ECG. In a study of healthy subjects, supratherapeutic doses of rilpivirine (75 mg once daily and 300 mg once daily) have been shown to prolong the QTc interval of the ECG (see section 5.1). Juluca should be used with caution when co-administered with a medicinal product with a known risk of Torsade de Pointes.

#### **4.6 Fertility, pregnancy and lactation**

##### Women of childbearing potential

Women of childbearing potential (WOCBP) should undergo pregnancy testing before initiation of Juluca. WOCBP who are taking Juluca should use effective contraception throughout treatment.

##### Pregnancy

There are limited data from the use of rilpivirine in pregnant women. Preliminary data from a surveillance study has suggested an increased incidence of neural tube defects (0.9%) in mothers exposed to dolutegravir at the time of conception compared with mothers exposed to non-dolutegravir containing regimens (0.1%).

The incidence of neural tube defects in the general population ranges from 0.5-1 case per 1,000 live births (0.05-0.1%). As neural tube defects occur within the first 4 weeks of foetal development (at which time the neural tubes are sealed) this potential risk would concern women exposed to dolutegravir at the time of conception and in early pregnancy. Due to the potential risk of neural tube defects, dolutegravir should not be used during the first trimester unless there is no alternative.

More than 1000 outcomes from second and third trimester exposure to dolutegravir in pregnant women indicate no evidence of increased risk of malformative and foeto/neonatal negative effects. However, as the mechanism by which dolutegravir may interfere in human pregnancy is unknown, the safety in use during the second and third trimester cannot be confirmed.

In animal reproductive toxicology studies with dolutegravir, no adverse development outcomes, including neural tube defects, were identified (see section 5.3). Dolutegravir was shown to cross the placenta in animals.

Animal studies with rilpivirine do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3).

Lower exposures of dolutegravir and rilpivirine were observed during pregnancy (see sections 4.2, 4.4, 5.1, 5.2).

The use of Juluca during pregnancy is not recommended.

#### Breast-feeding

It is unknown whether dolutegravir or rilpivirine are excreted in human milk. Available toxicological data in animals has shown excretion of dolutegravir and rilpivirine in milk. In lactating rats that received a single oral dose of dolutegravir 50 mg/kg at 10 days postpartum, dolutegravir was detected in milk at concentrations typically higher than blood. It is recommended that HIV infected women do not breast-feed their infants under any circumstances in order to avoid transmission of HIV.

#### Fertility

There are no data on the effects of dolutegravir or rilpivirine on human male or female fertility. Animal studies indicate no clinically relevant effects on male or female fertility (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Patients should be informed that fatigue, dizziness and somnolence have been reported during treatment with the components of Juluca. The clinical status of the patient and the adverse reaction profile of Juluca should be borne in mind when considering the patient's ability to drive or operate machinery.

### **4.8 Undesirable effects**

#### Summary of the safety profile

Clinical safety data with Juluca is limited. The most frequently reported adverse reactions considered possibly or probably related to the combined administration of dolutegravir plus rilpivirine in 513 HIV-1 infected subjects in the Phase III clinical trials (see section 5.1), were diarrhoea (2%) and headache (2%).

The most severe adverse reaction, possibly related to the treatment with dolutegravir (from pooled from Phase IIb and Phase III clinical studies), seen in an individual patient, was a hypersensitivity reaction that included rash and severe liver effects (see section 4.4).

#### Tabulated list of adverse reactions

The adverse reactions considered at least possibly related to treatment with the components of Juluca from clinical studies and post-marketing experience are listed in Table 2 by body system, organ class and frequency. Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), not known (cannot be estimated from the available data).

**Table 2: Tabulated summary of adverse reactions to Juluca based on clinical study and post-marketing experience with Juluca and its individual components**

System Organ Class (SOC)	Frequency category*	Adverse drug reactions
Blood and lymphatic systems disorders:	common	decreased white blood cell count decreased haemoglobin decreased platelet count
Immune system disorders	uncommon	hypersensitivity (see section 4.4)
	not known	immune reconstitution syndrome
Metabolism and nutrition disorders	very common	increased total cholesterol (fasted) increased LDL cholesterol (fasted)
	common	decreased appetite increased triglycerides (fasted)
Psychiatric disorders	very common	insomnia
	common	abnormal dreams depression sleep disorders depressed mood anxiety
	uncommon	suicidal ideation or suicide attempt (particularly in patients with a pre-existing history of depression or psychiatric illness)
	very common	headache dizziness
Nervous system disorders	common	somnolence
	very common	nausea increased pancreatic amylase diarrhoea
Gastrointestinal disorders	very common	

	common	abdominal pain vomiting flatulence increased lipase abdominal discomfort upper abdominal pain dry mouth
Hepatobiliary disorders	very common	increased transaminases (alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST) elevations)
	common	increased bilirubin
	uncommon	hepatitis
	rare	acute hepatic failure**
Skin and subcutaneous tissue disorders	common	rash pruritus
Musculoskeletal and connective tissue disorders	uncommon	arthralgia myalgia
General disorders and administration site conditions	common	fatigue
Investigations	common	creatinine phosphokinase (CPK) elevations

\* Frequencies are assigned based on the maximum frequencies observed in the pooled SWORD studies or studies with the individual components

\*\* This adverse reaction was identified through post-marketing surveillance for dolutegravir in combination with other ARVs. The frequency category of rare was estimated based on post-marketing reports.

#### Description of selected adverse reactions

##### *Changes in laboratory biochemistries*

Dolutegravir or rilpivirine have been associated with increases in serum creatinine occurring in the first week of treatment when administered with other antiretroviral medicinal products. Increases in serum creatinine occurred within the first four weeks of treatment with Juluca and remained stable through 48 weeks. A mean change from baseline of 8.22 µmol/L (range -26.5 to 51.2 µmol/L) was observed after 48 weeks treatment. These changes are related to inhibition of active transport, and are not considered to be clinically relevant as they do not reflect a change in glomerular filtration rate.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

#### **4.9 Overdose**

No specific symptoms or signs have been identified following acute overdose with dolutegravir or rilpivirine apart from those listed as adverse reactions.

Further management should be as clinically indicated or as recommended by the national poisons centre, where available. There is no specific treatment for an overdose of Juluca. If overdose occurs, the patient should be treated supportively with appropriate monitoring, including monitoring of vital signs and ECG (QT interval), as necessary. As dolutegravir and rilpivirine are highly bound to plasma proteins, dialysis is unlikely to result in significant removal of the active substances.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antivirals for systemic use, antivirals for treatment of HIV infections, combinations. ATC code: J05AR21

#### Mechanism of action

Dolutegravir inhibits HIV integrase by binding to the integrase active site and blocking the strand transfer step of retroviral Deoxyribonucleic acid (DNA) integration which is essential for the HIV replication cycle.

Rilpivirine is a diarylpyrimidine non-nucleoside reverse transcriptase inhibitor (NNRTI) of HIV-1. Rilpivirine activity is mediated by non-competitive inhibition of HIV-1 reverse transcriptase (RT). Rilpivirine does not inhibit the human cellular DNA polymerases  $\alpha$ ,  $\beta$  and  $\gamma$ .

#### Pharmacodynamic effects

##### *Antiviral activity in cell culture*

The IC<sub>50</sub> for dolutegravir against various laboratory strains using PBMC was 0.5 nM, and when using MT-4 cells it ranged from 0.7-2 nM. Similar IC<sub>50</sub>s were seen for clinical isolates without any major difference between subtypes; in a panel of 24 HIV-1 isolates of clades A, B, C, D, E, F and G and group O the mean IC<sub>50</sub> value was 0.2 nM (range 0.02-2.14). The mean IC<sub>50</sub> for 3 HIV-2 isolates was 0.18 nM (range 0.09-0.61).

Rilpivirine exhibited activity against laboratory strains of wild-type HIV-1 in an acutely infected T-cell line with a median IC<sub>50</sub> value for HIV-1/IIIB of 0.73 nM (0.27 ng/mL). Rilpivirine demonstrated limited *in vitro* activity against HIV-2 with IC<sub>50</sub> values ranging from 2,510 to 10,830 nM.

Rilpivirine also demonstrated antiviral activity against a broad panel of HIV-1 group M (clades A, B, C, D, F, G, H) primary isolates with IC<sub>50</sub> values ranging from 0.07 to 1.01 nM and group O primary isolates with EC<sub>50</sub> values ranging from 2.88 to 8.45 nM.

##### *Effect of human serum and serum proteins*

In 100% human serum, the dolutegravir mean protein fold shift was 75 fold, resulting in protein adjusted IC<sub>90</sub> of 0.064  $\mu$ g/mL.

A reduction in the antiviral activity of rilpivirine was observed in the presence of 1 mg/mL alpha-1-acid glycoprotein, 45 mg/mL human serum albumin, and 50% human serum as demonstrated by median IC<sub>50</sub> rates of 1.8, 39.2 and 18.5, respectively.

#### Resistance

##### *Resistance in vitro*

Serial passage is used to study resistance evolution *in vitro*. For dolutegravir, when using the laboratory strain HIV-1 IIIB during passage over 112 days, mutations selected appeared slowly, with substitutions at

positions S153Y and F; these mutations were not selected in patients treated with dolutegravir in the clinical studies. Using strain NL432, integrase mutations E92Q (fold change [FC] 3) and G193E (FC 3) were selected. These mutations have been selected in patients with pre-existing raltegravir resistance and who were then treated with dolutegravir (listed as secondary mutations for dolutegravir).

In further selection experiments using clinical isolates of subtype B, mutation R263K was seen in all five isolates (after 20 weeks and onwards). In subtype C (n=2) and A/G (n=2) isolates the integrase substitution R263K was selected in one isolate, and G118R in two isolates. R263K was reported from two individual patients with subtype B and subtype C in the Phase III clinical program for ART experienced, INI naive subjects, but without effects on dolutegravir susceptibility *in vitro*. G118R lowers the susceptibility to dolutegravir in site directed mutants (FC 10), but was not detected in patients receiving dolutegravir in the Phase III program.

Primary mutations for raltegravir/elvitegravir (Q148H/R/K, N155H, Y143R/H/C, E92Q, T66I) do not affect the *in vitro* susceptibility of dolutegravir as single mutations. When mutations listed as secondary integrase inhibitor associated mutations (for raltegravir/elvitegravir) are added to primary mutations (excluding at Q148) in experiments with site directed mutants, dolutegravir susceptibility remains at or near wildtype level. In the case of the Q148-mutation viruses, increasing dolutegravir FC is seen as the number of secondary mutations increase. The effect of the Q148-based mutations (H/R/K) was also consistent with *in vitro* passage experiments with site directed mutants. In serial passage with strain NL432-based site directed mutants at N155H or E92Q, no further selection of resistance was seen (FC unchanged around 1). In contrast, starting passage with mutants with mutation Q148H (FC 1), a variety of raltegravir associated secondary mutations accumulated with a consequent increase of FC to values >10.

A clinically relevant phenotypic cut-off value (FC vs wild type virus) has not been determined; genotypic resistance was a better predictor for outcome.

Rilpivirine-resistant strains were selected in cell culture starting from wild type HIV-1 of different origins and clades as well as NNRTI-resistant HIV-1. The most commonly observed amino acid substitutions that emerged included: L100I, K101E, V108I, E138K, V179F, Y181C, H221Y, F227C and M230I. Resistance to rilpivirine was considered present when FC in EC<sub>50</sub> value was above the biological cut-off (BCO) of the assay.

#### *Resistance in vivo*

Two subjects across the pooled SWORD-1 (201636) and SWORD-2 (201637) studies from each treatment arm experienced confirmed virologic failure leading to withdrawal at any time through Week 48. NNRTI resistance associated substitution K101K/E mixture with no decreased susceptibility to rilpivirine (FC=1.2) was observed in one subject with identified adherence issues that received dolutegravir plus rilpivirine. No integrase resistance was observed. This patient's viral load was 1,059,771 copies/mL at the suspected virologic withdrawal visit and was <50 copies/mL at the withdrawal visit. No treatment emergent resistance-associated substitutions were observed for the other three subjects who experienced confirmed virologic failure.

In previously untreated patients receiving dolutegravir + 2 NRTIs in Phase IIb and Phase III, no development of resistance to the integrase class, or to the NRTI class was seen (n=876, follow-up of 48-96 weeks).

In patients with prior failed therapies, but naïve to the integrase class (SAILING study), integrase inhibitor substitutions were observed in 4/354 patients (follow-up 48 weeks) treated with dolutegravir, which was given in combination with an investigator selected background regimen (BR). Of these four, two subjects had a unique R263K integrase substitution, with a maximum FC of 1.93, one subject had a polymorphic V151V/I integrase substitution, with maximum FC of 0.92, and one subject had pre-existing integrase mutations and is assumed to have been integrase inhibitor experienced or infected with integrase inhibitor resistant virus by transmission. The R263K mutation was also selected *in vitro* (see above).

From rilpivirine Phase III trials, in the week 48 pooled resistance analysis conducted with previously untreated patients, 62 (of a total of 72) virologic failures in the rilpivirine arm had resistance data at baseline

and time of failure. In this analysis, the resistance-associated mutations (RAMs) associated with NNRTI resistance that developed in at least 2 rilpivirine virologic failures were: V90I, K101E, E138K, E138Q, V179I, Y181C, V189I, H221Y, and F227C. In the trials, the presence of the mutations V90I and V189I, at baseline, did not affect response. The E138K substitution emerged most frequently during rilpivirine treatment, commonly in combination with the M184I substitution. In the week 48 analysis, 31 out of 62 of rilpivirine virologic failures had concomitant NNRTI and NRTI RAMs; 17 of those 31 had the combination of E138K and M184I. The most common mutations were the same in the week 48 and week 96 analyses. From the week 48 to the week 96 analysis, 24 (3.5%) and 14 (2.1%) additional virologic failures occurred in the rilpivirine and efavirenz arm, respectively.

### Cross-resistance

#### *Site-directed INI mutant virus*

Dolutegravir activity was determined against a panel of 60 INI-resistant site-directed mutant HIV-1 viruses (28 with single substitutions and 32 with 2 or more substitutions). The single INI-resistance substitutions T66K, I151L, and S153Y conferred a greater than 2-fold decrease in dolutegravir susceptibility (range: 2.3-fold to 3.6-fold from reference). Combinations of multiple substitutions T66K/L74M, E92Q/N155H, G140C/Q148R, G140S/Q148H, R or K, Q148R/N155H, T97A/G140S/Q148, and substitutions at E138/G140/Q148 showed a greater than 2-fold decrease in dolutegravir susceptibility (range: 2.5-fold to 21-fold from reference).

#### *Site-directed NNRTI mutant virus*

In a panel of 67 HIV-1 recombinant laboratory strains with one amino acid substitution at RT positions associated with NNRTI resistance, including the most commonly found K103N and Y181C, rilpivirine showed antiviral activity ( $FC \leq BCO$ ) against 64 (96%) of these strains. The single amino acid substitutions associated with a loss of susceptibility to rilpivirine were: K101P, Y181I and Y181V. The K103N substitution did not result in reduced susceptibility to rilpivirine by itself, but the combination of K103N and L100I resulted in a 7-fold reduced susceptibility to rilpivirine.

Considering all of the available *in vitro* and *in vivo* data, the following amino acid substitutions, when present at baseline, are likely to affect the activity of rilpivirine: K101E, K101P, E138A, E138G, E138K, E138R, E138Q, V179L, Y181C, Y181I, Y181V, Y188L, H221Y, F227C, M230I or M230L.

#### *Recombinant clinical isolates*

Seven hundred and five raltegravir resistant isolates from raltegravir experienced patients were analysed for susceptibility to dolutegravir. Dolutegravir had a <10 FC against 94% of the 705 clinical isolates.

Rilpivirine retained sensitivity ( $FC \leq BCO$ ) against 62% of 4786 HIV-1 recombinant clinical isolates resistant to efavirenz and/or nevirapine.

#### *Previously untreated HIV-1 infected adult patients*

In a Week 96 pooled analyses of virologic failures with baseline viral load  $\leq 100,000$  copies/mL and resistance to rilpivirine ( $n = 5$ ), subjects had cross-resistance to efavirenz ( $n = 3$ ), etravirine ( $n = 4$ ), and nevirapine ( $n = 1$ ).

### Effects on electrocardiogram

The effect of rilpivirine at the recommended dose of 25 mg once daily on the QTcF interval was evaluated in a randomised, placebo and active (moxifloxacin 400 mg once daily) controlled crossover study in 60 healthy adults, with 13 measurements over 24 hours at steady-state. Rilpivirine at the recommended dose of 25 mg once daily is not associated with a clinically relevant effect on QTc.

When supratherapeutic doses of 75 mg once daily and 300 mg once daily of rilpivirine were studied in healthy adults, the maximum mean time-matched (95% upper confidence bound) differences in QTcF interval from placebo after baseline correction were 10.7 (15.3) and 23.3 (28.4) ms, respectively. Steady-state administration of rilpivirine 75 mg once daily and 300 mg once daily resulted in a mean  $C_{max}$

approximately 2.6-fold and 6.7-fold, respectively, higher than the mean steady-state C<sub>max</sub> observed with the recommended 25 mg once daily dose of rilpivirine (see section 4.4).

No relevant effects were seen with dolutegravir on the QTc interval, with doses exceeding the clinical dose by approximately three fold.

#### Clinical efficacy and safety

The efficacy and safety of switching from an antiretroviral regimen (containing 2 NRTIs plus either an INI, an NNRTI, or a PI) to a dual regimen of dolutegravir 50 mg and rilpivirine 25 mg was evaluated in 2 identical 48-week, randomised, open-label, multicentre, parallel-group, non-inferiority studies SWORD-1 (201636) and SWORD-2 (201637). Subjects were enrolled if they were on their first or second antiretroviral regimen with no history of virological failure, had no suspected or known resistance to any antiretroviral and had been stably suppressed (HIV-1 RNA < 50 copies/mL) for at least 6 months prior to screening. Subjects were randomised 1:1 to continue their CAR or be switched to a two-drug regimen dolutegravir plus rilpivirine administered once daily. The primary efficacy endpoint for the SWORD studies was the proportion of subjects with plasma HIV-1 RNA <50 copies/mL at Week 48 (Snapshot algorithm for the ITT-E population).

At baseline, in the pooled analysis, characteristics were similar between treatment arms with the median age of subjects of 43 years (28%, 50 years of age or older; 3%, 65 years of age or older), 22% female, 20% non-white and 77% were CDC Class A. Median CD+cell count was about 600 cells per mm<sup>3</sup> with 11% having CD4+ cell count less than 350 cells per mm<sup>3</sup>. In the pooled analysis, 54%, 26%, and 20% of subjects were receiving an NNRTI, PI, or INI (respectively) as their baseline third treatment agent class prior to randomisation.

The pooled primary analysis demonstrated that dolutegravir plus rilpivirine is non-inferior to CAR, with 95% of subjects in both arms achieving the primary endpoint of <50 copies/mL plasma HIV-1 RNA at Week 48 based on the Snapshot algorithm (Table 3).

The primary endpoint and other outcomes (including outcomes by key baseline covariates) for the pooled SWORD-1 and SWORD-2 studies are shown in Table 3.

**Table 3: Virologic Outcomes of Randomized Treatment at Week 48 (Snapshot algorithm)**

	SWORD-1 and SWORD-2 Pooled Data***	
	DTG + RPV N=513 n (%)	CAR N=511 n (%)
<b>HIV-1 RNA &lt;50 copies/mL</b>	486 (95%)	485 (95%)
<b>Treatment Difference*</b>	-0.2 (-3.0, 2.5)	
<b>Virologic non response**</b>		
<u>Reasons</u>		
Data in window not <50 copies/mL	3 (<1%)	6 (1%)
Discontinued for lack of efficacy	0	2 (<1%)
Discontinued for other reasons while not <50 copies/mL	2 (<1%)	2 (<1%)
Change in ART	1 (<1%)	1 (<1%)
Change in ART	0	1 (<1%)
<b>No virologic data at Week 48 window</b>	24 (5%)	20 (4%)
<u>Reasons</u>		
Discontinued study/study drug due to adverse event or death	17 (3%)	3 (<1%)
Discontinued study/study drug for other reasons	7 (1%)	16 (3%)
Missing data during window but on study	0	1 (<1%)
<b>HIV-1 RNA &lt;50 copies/mL by baseline covariates</b>		

	n/N (%)	n/N (%)
<b>Baseline CD4+ (cells/ mm<sup>3</sup>)</b>		
<350	51 / 58 (88%)	46 / 52 (88%)
≥350	435 / 455 (96%)	439 / 459 (96%)
<b>Baseline Third Treatment Agent Class</b>		
INI	99 / 105 (94%)	92 / 97 (95%)
NNRTI	263 / 275 (96%)	265 / 278 (95%)
PI	124 / 133 (93%)	128 / 136 (94%)
<b>Gender</b>		
Male	375 / 393 (95%)	387 / 403 (96%)
Female	111 / 120 (93%)	98 / 108 (91%)
<b>Race</b>		
White	395 / 421 (94%)	380 / 400 (95%)
African-America/African Heritage/Other	91 / 92 (99%)	105 / 111 (95%)
<b>Age (years)</b>		
<50	350 / 366 (96%)	348 / 369 (94%)
≥50	136 / 147 (93%)	137 / 142 (96%)

\* Adjusted for baseline stratification factors and assessed using a non-inferiority margin of - 8%.

\*\* Non-inferiority of dolutegravir plus rilpivirine to CAR, in the proportion of subjects classified as virologic non-responders, was demonstrated using a non-inferiority margin of 4%. Adjusted difference (95% CI) -0.6 (-1.7, 0.6).

\*\*\* The results of the pooled analysis are in line with those of the individual studies, for which differences in proportions meeting the primary endpoint of <50 copies/mL plasma HIV-1 RNA at Week 48 (based on the Snapshot algorithm) for DTG+RPV versus CAR were -0.6 (95% CI: -4.3; 3.0) for SWORD-1 and 0.2 (95% CI: -3.9; 4.2) for SWORD-2 with a preset non-inferiority margin of -10%.

N = Number of subjects in each treatment arm  
CAR = current antiretroviral regimen; DTG+RPV = dolutegravir plus rilpivirine;  
INI = Integrase inhibitor; NNRTI = Non-nucleoside reverse transcriptase inhibitor;  
PI = Protease Inhibitor

#### *Effects on bone*

In a DEXA substudy mean bone mineral density (BMD) increased from Baseline to Week 48 in subjects who switched to dolutegravir plus rilpivirine (1.34% total hip and 1.46% lumbar spine) compared with those who continued on treatment with a TDF-containing antiretroviral regimen (0.05% total hip and 0.15% lumbar spine). Any beneficial effect on fracture rate was not studied.

#### Pregnancy

No efficacy and safety data are available for the combination of dolutegravir and rilpivirine in pregnancy. Rilpivirine in combination with a background regimen was evaluated in a clinical trial of 19 pregnant women during the second and third trimesters, and postpartum. The pharmacokinetic data demonstrate that total exposure (AUC) to rilpivirine as a part of an antiretroviral regimen was approximately 30% lower during pregnancy compared with postpartum (6-12 weeks). Of the 12 subjects that completed the study, 10 subjects were suppressed at the end of the study; in the other 2 subjects an increase in viral load was observed postpartum, for 1 subject due to suspected suboptimal adherence. No mother to child transmission occurred in all 10 infants born to the mothers who completed the trial and for whom the HIV status was available. There were no new safety findings compared with the known safety profile of rilpivirine in HIV-1 infected adults.

In limited data from small numbers of women who received dolutegravir 50 mg once daily in combination with a background regimen, the total exposure (AUC) to dolutegravir was 37% lower during the 2nd trimester of pregnancy, and 29% lower during the 3rd trimester of pregnancy, compared with postpartum (6-12 weeks). Of the 29 subjects that completed the study, 27 subjects were suppressed at the end of the study.

No mother to child transmission was identified. While 24 infants were confirmed to be uninfected, 5 were indeterminate due to incomplete testing, see sections 4.2, 4.4 and 5.2.

#### Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Juluca in one or more subsets of the paediatric population in the treatment of HIV infection.

### **5.2 Pharmacokinetic properties**

Juluca is bioequivalent to a dolutegravir 50 mg tablet and a rilpivirine 25 mg tablet administered together with a meal.

Dolutegravir pharmacokinetics are similar between healthy and HIV-infected subjects. The PK variability of dolutegravir is low to moderate. In Phase I studies in healthy subjects, between-subject CVb% for AUC and C<sub>max</sub> ranged from ~20 to 40% and C<sub>T</sub> from 30 to 65% across studies. The between-subject PK variability of dolutegravir was higher in HIV-infected subjects than healthy subjects. Within-subject variability (CVw%) is lower than between-subject variability.

The pharmacokinetic properties of rilpivirine have been evaluated in adult healthy subjects and in adult antiretroviral treatment-naïve HIV-1 infected patients. Systemic exposure to rilpivirine was generally lower in HIV-1 infected patients than in healthy subjects.

#### Absorption

Dolutegravir is rapidly absorbed following oral administration, with median T<sub>max</sub> at 2 to 3 hours post dose for tablet formulation. After oral administration, the maximum plasma concentration of rilpivirine is generally achieved within 4-5 hours.

Juluca must be taken with a meal to obtain optimal absorption of rilpivirine (see section 4.2). When Juluca was taken with a meal, the absorption of both dolutegravir and rilpivirine was increased. Moderate and high fat meals increased the dolutegravir AUC(0-∞) by approximately 87% and C<sub>max</sub> by approximately 75%. Rilpivirine AUC(0-∞) was increased by 57% and 72% and C<sub>max</sub> by 89% and 117%, with moderate and high fat meals respectively, compared to fasted conditions. Taking Juluca in fasted condition or with only a protein-rich nutritional drink may result in decreased plasma concentrations of rilpivirine, which could potentially reduce the therapeutic effect of Juluca.

The absolute bioavailability of dolutegravir or rilpivirine has not been established.

#### Distribution

Dolutegravir is highly bound (>99%) to human plasma proteins based on *in vitro* data. The apparent volume of distribution is 17 L to 20 L in HIV-infected patients, based on a population pharmacokinetic analysis. Binding of dolutegravir to plasma proteins is independent of dolutegravir concentration. Total blood and plasma drug-related radioactivity concentration ratios averaged between 0.441 to 0.535, indicating minimal association of radioactivity with blood cellular components. The unbound fraction of dolutegravir in plasma is increased at low levels of serum albumin (<35 g/L) as seen in subjects with moderate hepatic impairment.

Dolutegravir is present in cerebrospinal fluid (CSF). In 13 treatment-naïve subjects on a stable dolutegravir plus abacavir/lamivudine regimen, dolutegravir concentration in CSF averaged 18 ng/mL (comparable to unbound plasma concentration, and above the IC<sub>50</sub>).

Dolutegravir is present in the female and male genital tract. AUC in cervicovaginal fluid, cervical tissue and vaginal tissue were 6-10% of those in corresponding plasma at steady state. AUC in semen was 7% and 17% in rectal tissue of those in corresponding plasma at steady state.

Rilpivirine is approximately 99.7% bound to plasma proteins *in vitro*, primarily to albumin. The distribution of rilpivirine into compartments other than plasma (e.g., cerebrospinal fluid, genital tract secretions) has not been evaluated in humans.

### Biotransformation

Dolutegravir is primarily metabolized through glucuronidation via UGT1A1 with a minor CYP3A component. Dolutegravir is the predominant circulating compound in plasma; renal elimination of unchanged active substance is low (< 1% of the dose). Fifty-three percent of total oral dose is excreted unchanged in the faeces. It is unknown if all or part of this is due to unabsorbed active substance or biliary excretion of the glucuronide conjugate, which can be further degraded to form the parent compound in the gut lumen. Thirty-two percent of the total oral dose is excreted in the urine, mainly represented by ether glucuronide of dolutegravir (18.9% of total dose), N-dealkylation metabolite (3.6% of total dose), and a metabolite formed by oxidation at the benzylic carbon (3.0% of total dose).

*In vitro* experiments indicate that rilpivirine primarily undergoes oxidative metabolism mediated by the CYP3A system.

### Drug interactions

*In vitro*, dolutegravir demonstrated no direct, or weak inhibition ( $IC_{50}>50\text{ }\mu\text{M}$ ) of the enzymes cytochrome P450 (CYP)1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 CYP3A, uridine diphosphate glucuronosyl transferase (UGT)1A1 or UGT2B7, or the transporters Pgp, BCRP, BSEP, OATP1B1, OATP1B3, OCT1, MATE2-K, MRP2 or MRP4. *In vitro*, dolutegravir did not induce CYP1A2, CYP2B6 or CYP3A4. Based on this data, dolutegravir is not expected to affect the pharmacokinetics of medicinal products that are substrates of major enzymes or transporters (see section 4.5).

*In vitro*, dolutegravir was not a substrate of human OATP 1B1, OATP 1B3 or OCT 1.

### Elimination

Dolutegravir has a terminal half-life of ~14 hours. The apparent oral clearance (CL/F) is approximately 1L/hr in HIV-infected patients based on a population pharmacokinetic analysis.

The terminal elimination half-life of rilpivirine is approximately 45 hours. After single dose oral administration of  $^{14}\text{C}$ -rilpivirine, on average 85% and 6.1% of the radioactivity could be retrieved in faeces and urine, respectively. In faeces, unchanged rilpivirine accounted for on average 25% of the administered dose. Only trace amounts of unchanged rilpivirine (< 1% of dose) were detected in urine.

### Special patient populations

#### *Children*

Neither Juluca nor the combination dolutegravir and rilpivirine as single entities have been studied in children. Dosing recommendations for paediatric patients cannot be made due to insufficient data (see section 4.2).

The pharmacokinetics of dolutegravir in 10 antiretroviral treatment-experienced HIV-1 infected adolescents (12 to <18 years of age and weighing  $\geq 40\text{ kg}$ ) showed that dolutegravir 50 mg once daily oral dosage resulted in dolutegravir exposure comparable to that observed in adults who received dolutegravir 50 mg orally once daily. The pharmacokinetics was evaluated in 11 children 6 to 12 years of age and showed that 25 mg once daily in patients weighing at least 20 kg and 35 mg once daily in patients weighing at least 30 kg resulted in dolutegravir exposure comparable to adults.

The pharmacokinetics of rilpivirine in 36 antiretroviral treatment-naïve HIV-1 infected adolescent subjects (12 to <18 years of age) receiving rilpivirine 25 mg once daily were comparable to those in treatment-naïve HIV-1 infected adults receiving rilpivirine 25 mg once daily. There was no impact of body weight on

rilpivirine pharmacokinetics in paediatric subjects in study C213 (33 to 93 kg), similar to what was observed in adults.

#### *Elderly*

Population pharmacokinetic analysis using data in HIV-1 infected adults showed that there was no clinically relevant effect of age on dolutegravir or rilpivirine exposures. Pharmacokinetic data in subjects >65 years old are very limited.

#### *Renal impairment*

Renal clearance of unchanged active substance is a minor pathway of elimination for dolutegravir. A study of the pharmacokinetics of dolutegravir was performed in subjects with severe renal impairment (CLcr <30 mL/min) and matched healthy controls. The exposure to dolutegravir was decreased by approximately 40% in subjects with severe renal impairment. The mechanism for the decrease is unknown. The pharmacokinetics of rilpivirine have not been studied in patients with renal insufficiency.

Renal elimination of rilpivirine is negligible. No dose adjustment is needed for patients with mild or moderate renal impairment. In patients with severe renal impairment or end-stage renal disease, Juluca should be used with caution, as rilpivirine plasma concentrations may be increased due to alteration of drug absorption, distribution and/or metabolism secondary to renal dysfunction. In patients with severe renal impairment or end-stage renal disease, the combination of Juluca with a strong CYP3A inhibitor should only be used if the benefit outweighs the risk. Juluca has not been studied in patients on dialysis. As dolutegravir and rilpivirine are highly bound to plasma proteins, it is unlikely that they will be significantly removed by haemodialysis or peritoneal dialysis (see section 4.2).

#### *Hepatic impairment*

Dolutegravir and rilpivirine are both primarily metabolized and eliminated by the liver. A single dose of 50 mg of dolutegravir was administered to 8 subjects with moderate hepatic impairment (Child-Pugh score B) and to 8 matched healthy adult controls. While the total dolutegravir concentration in plasma was similar, a 1.5- to 2-fold increase in unbound exposure to dolutegravir was observed in subjects with moderate hepatic impairment compared to healthy controls.

In a rilpivirine study comparing 8 patients with mild hepatic impairment (Child-Pugh score A) to 8 matched controls, and 8 patients with moderate hepatic impairment (Child-Pugh score B) to 8 matched controls, the multiple dose exposure of rilpivirine was 47% higher in patients with mild hepatic impairment and 5% higher in patients with moderate hepatic impairment. However, it may not be excluded that the pharmacologically active, unbound, rilpivirine exposure is significantly increased in moderate hepatic impairment.

No dosage adjustment is considered necessary for patients with mild to moderate hepatic impairment (Child-Pugh score A or B). Juluca should be used with caution in patients with moderate hepatic impairment. The effect of severe hepatic impairment (Child-Pugh score C) on the pharmacokinetics of dolutegravir or rilpivirine has not been studied, therefore Juluca is not recommended in these patients.

#### *Gender*

Population pharmacokinetic analyses from studies with the individual components revealed that gender had no clinically relevant effect on the pharmacokinetics of dolutegravir or rilpivirine.

#### *Race*

No clinically important pharmacokinetic differences of dolutegravir or rilpivirine due to race have been identified.

#### *Co-infection with Hepatitis B or C*

Population pharmacokinetic analysis indicated that hepatitis C virus co-infection had no clinically relevant effect on the exposure to dolutegravir or rilpivirine. Subjects with hepatitis B co-infection or hepatitis C infection in need of anti-HCV therapy were excluded from studies with the dual combination of dolutegravir and rilpivirine.

#### *Pregnancy and postpartum*

No pharmacokinetic data are available for the combination of dolutegravir and rilpivirine in pregnancy. In limited data from small numbers of women in study IMPAACT P1026 who received dolutegravir 50 mg once daily during the 2nd trimester of pregnancy, mean intra-individual values for total dolutegravir  $C_{\max}$ ,  $AUC_{24h}$  and  $C_{24h}$  values were, respectively, 26%, 37% and 51% lower as compared to postpartum; during the 3rd trimester of pregnancy,  $C_{\max}$ ,  $AUC_{24h}$  and  $C_{\min}$  values were, respectively, 25%, 29% and 34% lower as compared to postpartum (see sections 4.2, 4.4. and 4.6).

In women receiving rilpivirine 25 mg once daily during the 2nd trimester of pregnancy, mean intra-individual values for total rilpivirine  $C_{\max}$ ,  $AUC_{24h}$  and  $C_{\min}$  values were, respectively, 21%, 29% and 35% lower as compared to postpartum; during the 3rd trimester of pregnancy,  $C_{\max}$ ,  $AUC_{24h}$  and  $C_{\min}$  values were, respectively, 20%, 31% and 42% lower as compared to postpartum (see sections 4.2, 4.4. and 4.6).

### **5.3 Preclinical safety data**

#### Carcinogenesis and mutagenesis

Dolutegravir was not mutagenic or clastogenic using *in vitro* tests in bacteria and cultured mammalian cells, and an *in vivo* rodent micronucleus assay. Dolutegravir was not carcinogenic in long term studies in the mouse and rat.

Rilpivirine has tested negative in the absence and presence of a metabolic activation system in the *in vitro* Ames reverse mutation assay and the *in vitro* clastogenicity mouse lymphoma assay. Rilpivirine did not induce chromosomal damage in the *in vivo* micronucleus test in mice. Carcinogenicity studies with rilpivirine in mice and rats revealed tumorigenic potential specific for these species, but are regarded as of no relevance for humans.

#### Reproductive toxicology studies

Dolutegravir did not affect male or female fertility in rats at doses up to 1000 mg/kg/day, the highest dose tested (33 times the 50 mg human clinical exposure based on AUC).

Oral administration of dolutegravir to pregnant rats at doses up to 1000 mg/kg daily from days 6 to 17 of gestation did not elicit maternal toxicity, developmental toxicity or teratogenicity (38 times the 50 mg human clinical exposure based on AUC).

Oral administration of dolutegravir to pregnant rabbits at doses up to 1000 mg/kg daily from days 6 to 18 of gestation did not elicit developmental toxicity or teratogenicity (0.56 times the 50 mg human clinical exposure based on AUC). In rabbits, maternal toxicity (decreased food consumption, scant/no faeces/urine, suppressed body weight gain) was observed at 1000 mg/kg (0.56 times the 50 mg human clinical exposure based on AUC).

Rilpivirine studies in animals have shown no evidence of relevant embryonic or foetal toxicity or an effect on reproductive function. There was no teratogenicity with rilpivirine in rats and rabbits. The exposures at the embryo-foetal No Observed Adverse Effects Levels (NOAELs) in rats and rabbits were respectively 15 and 70 times higher than the exposure in humans at the recommended dose of 25 mg once daily.

#### Repeated dose toxicity

The effect of prolonged daily treatment with high doses of dolutegravir has been evaluated in repeat oral dose toxicity studies in rats (up to 26 weeks) and in monkeys (up to 38 weeks). The primary effect of dolutegravir was gastrointestinal intolerance or irritation in rats and monkeys at doses that produce systemic exposures approximately 30 and 1.2 times the 50 mg human clinical exposure based on AUC, respectively. Because gastrointestinal (GI) intolerance is considered to be due to local active substance administration,

mg/kg or mg/m<sup>2</sup> metrics are appropriate determinates of safety cover for this toxicity. GI intolerance in monkeys occurred at 30 times the human mg/kg equivalent dose (based on a 50 kg human), and 11 times the human mg/m<sup>2</sup> equivalent dose for a clinical dose of 50 mg.

Liver toxicity associated with liver enzyme induction was observed in rodents following rilpivirine administration. In dogs, cholestasis-like effects were noted.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Mannitol (E421)  
Magnesium stearate  
Microcrystalline cellulose  
Povidone (K29/32)  
Sodium starch glycolate  
Sodium stearyl fumarate  
Lactose monohydrate  
Croscarmellose sodium  
Povidone (K30)  
Polysorbate 20  
Silicified microcrystalline cellulose

#### Tablet coating

Polyvinyl alcohol- part hydrolysed  
Titanium dioxide (E171)  
Macrogol  
Talc  
Iron oxide yellow (E172)  
Iron oxide red (E172)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

3 years.

### **6.4 Special precautions for storage**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

This medicinal product does not require any special temperature storage conditions.

### **6.5 Nature and contents of container**

White HDPE (high density polyethylene) bottles closed with polypropylene child-resistant closures, with a polyethylene faced induction heat seal liner. Each pack consists of one bottle containing 30 film-coated tablets and a desiccant.

Multipacks containing 90 (3 packs of 30) film-coated tablets. Each pack of 30 film-coated tablets contains a desiccant.

Not all pack sizes may be marketed.

#### **6.6 Special precautions for disposal**

No special requirements for disposal.

### **7. MARKETING AUTHORISATION HOLDER**

ViiV Healthcare BV  
Huis ter Heideweg 62  
3705 LZ Zeist  
Netherlands

### **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1282/001  
EU/1/18/1282/002

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 16 May 2018

### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

Glaxo Wellcome, S.A.  
Avda. Extremadura, 3  
09400 Aranda De Duero  
Burgos  
Spain

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- Periodic safety update reports**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder shall submit the first periodic safety update report for this product within 6 months following authorisation.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- Risk Management Plan (RMP)**

The MAH shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON (INDIVIDUAL PACKS ONLY)****1. NAME OF THE MEDICINAL PRODUCT**

Juluca 50 mg/25 mg film-coated tablets  
dolutegravir/rilpivirine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains dolutegravir sodium equivalent to 50 mg dolutegravir and rilpivirine hydrochloride equivalent to 25 mg rilpivirine.

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate.  
See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

30 tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR  
WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF  
APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare BV  
Huis ter Heideweg 62  
3705 LZ Zeist  
Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1282/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

juluca

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****OUTER CARTON (MULITPACKS ONLY – WITH BLUE BOX)****1. NAME OF THE MEDICINAL PRODUCT**

Juluca 50 mg/25 mg film-coated tablets  
dolutegravir/rilpivirine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains dolutegravir sodium equivalent to 50 mg dolutegravir and rilpivirine hydrochloride equivalent to 25 mg rilpivirine.

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate.  
See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Multipack: 90 (3 packs of 30) tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR  
WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF  
APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare BV  
Huis ter Heideweg 62  
3705 LZ Zeist  
Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1282/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

juluca

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC:  
SN:  
NN:

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING****INTERMEDIATE CARTON (WITHOUT BLUE BOX – COMPONENT OF MULTIPACK)****1. NAME OF THE MEDICINAL PRODUCT**

Juluca 50 mg/25 mg film-coated tablets  
dolutegravir/rilpivirine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains dolutegravir sodium equivalent to 50 mg dolutegravir and rilpivirine hydrochloride equivalent to 25 mg rilpivirine.

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate.  
See package leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

30 tablets. Component of a multipack, can't be sold separately.

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.  
Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR  
WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF  
APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare BV  
Huis ter Heideweg 62  
3705 LZ Zeist  
Netherlands

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1282/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

juluca

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING****BOTTLE LABEL****1. NAME OF THE MEDICINAL PRODUCT**

Juluca 50 mg/25 mg film-coated tablets  
dolutegravir/rilpivirine

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains dolutegravir sodium equivalent to 50 mg dolutegravir and rilpivirine hydrochloride equivalent to 25 mg rilpivirine.

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate.  
See leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

30 tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY****8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR  
WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF  
APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

ViiV Healthcare BV

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/18/1282/001

EU/1/18/1282/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

## **B. PACKAGE LEAFLET**

## **Package leaflet: Information for the patient**

### **Juluca 50 mg/25 mg film-coated tablets dolutegravir/rilpivirine**

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### **Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Juluca is and what it is used for
2. What you need to know before you take Juluca
3. How to take Juluca
4. Possible side effects
5. How to store Juluca
6. Contents of the pack and other information

#### **1. What Juluca is and what it is used for**

Juluca is a medicine that contains two active ingredients used to treat human immunodeficiency virus (HIV) infection: dolutegravir and rilpivirine. Dolutegravir belongs to a group of anti-retroviral medicines called *integrase inhibitors (INIs)*, and rilpivirine belongs to a group of anti-retroviral medicines called *non-nucleoside reverse transcriptase inhibitors (NNRTIs)*.

Juluca is used to treat HIV in adults aged 18 years and over who are taking other antiretroviral medicines and whose HIV-1 infection is under control for at least 6 months. Juluca may replace your current antiretroviral medicines.

Juluca keeps the amount of HIV virus in your body at a low level. This helps maintain the number of CD4 cells in your blood. CD4 cells are a type of white blood cells that are important in helping your body to fight infection.

#### **2. What you need to know before you take Juluca**

##### **Do not take Juluca:**

- if you are allergic to dolutegravir or rilpivirine or any of the other ingredients of this medicine (listed in section 6).

**Do not take Juluca if you are taking any of the following medicines** as they may affect the way Juluca works:

- fampridine (also known as dalfampridine; used in multiple sclerosis)
- carbamazepine, oxcarbazepine, phenobarbital, phenytoin (medicines to treat epilepsy and to prevent fits)
- rifampicin, rifapentine (medicines to treat some bacterial infections such as tuberculosis)

- omeprazole, esomeprazole, lansoprazole, pantoprazole, rabeprazole, (medicines to prevent and treat stomach ulcers, heartburn or acid reflux disease)
- dexamethasone (a corticosteroid used in many conditions such as inflammation and allergic reactions) when taken by mouth or injected, except as a single dose treatment
- products that contain St John's wort (*Hypericum perforatum*) (a herbal product used for depression).

If you are taking any of the above, ask your doctor about alternatives.

### **Warnings and precautions**

Talk to your doctor or, pharmacist before taking Juluca.

#### Allergic reactions

Juluca contains dolutegravir. Dolutegravir can cause a serious allergic reaction known as a hypersensitivity reaction. You need to know about important signs and symptoms to look out for while you're taking Juluca.

→ **Read the information** 'Allergic reactions' in section 4 of this leaflet.

#### Liver problems including hepatitis B and/or C

Tell your doctor if you have or have had **problems with your liver**, including hepatitis B and/or C. Your doctor may evaluate how severe your liver disease is before deciding if you can take Juluca.

#### Look out for important symptoms

Some people taking medicines for HIV infection develop other conditions, which can be serious. These include:

- symptoms of infections and inflammation
- joint pain, stiffness and bone problems.

You need to know about important signs and symptoms to look out for while you're taking Juluca.

→ **Read the information** 'Other possible side effects' in section 4 of this leaflet.

#### Protect other people

HIV infection is spread by sexual contact with someone who has the infection, or by transfer of infected blood (for example, by sharing injection needles). You can still pass on HIV when taking this medicine, although the risk is lowered by effective antiretroviral therapy. Discuss with your doctor the precautions needed to avoid infecting other people.

### **Children and adolescents**

This medicine is not for use in children or adolescents less than 18 years of age, because it has not been studied in these patients.

### **Other medicines and Juluca**

Tell your doctor if you are taking, have recently taken or might take any other medicines.

**Juluca must not be taken** with some other medicines (see 'Do not take Juluca' earlier in section 2).

Some medicines can affect how Juluca works, or make it more likely that you will have side effects. Juluca can also affect how some other medicines work.

**Tell your doctor** if you are taking any of the medicines *in the following list*:

- metformin, to treat **diabetes**
- medicines that may cause a life threatening irregular heartbeat (*Torsade de Pointes*). As a number of different medicines can cause this condition, you should ask your doctor or pharmacist if you are not sure
- medicines called **antacids**, to treat **indigestion** and **heartburn**. **Do not take an antacid** during the 6 hours before you take Juluca, or for at least 4 hours after you take it (see also section 3, 'How to take Juluca')

- **calcium supplements, iron supplements and multivitamins** must be taken at the same time as Juluca with a meal. If you can't take these supplements at the same time as Juluca, **do not take a calcium supplement, iron supplement or multivitamin** during the 6 hours before you take Juluca, or for at least 4 hours after you take it (see also section 3, 'How to take Juluca')
- medicines called **H<sub>2</sub> receptor antagonists** (for example cimetidine, famotidine, nizatidine, ranitidine) to treat stomach or **intestinal ulcers** or used to **relieve heartburn due to acid reflux**. **Do not take these medicines** during the 12 hours before you take Juluca, or for at least 4 hours after you take it (see also section 3, 'How to take Juluca')
- any medicines to treat **HIV infection**
- rifabutin, to treat tuberculosis (TB) and other **bacterial infections**. If you take rifabutin, your doctor may need to give you an additional dose of rilpivirine to treat your HIV infection (see section 3, 'How to take Juluca')
- artemether/lumefantrine used to prevent you catching **malaria**
- clarithromycin and erythromycin, to treat **bacterial infections**
- methadone, used in the treatment of opioid dependence
- dabigatran etexilate, used to treat or prevent **blood clots**.

→ **Tell your doctor or pharmacist** if you are taking any of these. Your doctor may decide that you need extra check ups.

### **Pregnancy**

If you are pregnant, think you may be pregnant, or if you are planning to have a baby:

→ **Use of Juluca is not recommended. Ask your doctor** for advice.

Taking Juluca at the time of becoming pregnant or during the first twelve weeks of pregnancy, may increase the risk of a type of birth defect, called neural tube defect, such as spina bifida (malformed spinal cord).

If you could get pregnant while receiving Juluca, you need to use a reliable method of barrier contraception (for example, a condom) with other methods of contraception including oral (pill) or other hormonal contraceptives (for example, implants, injection), to prevent pregnancy.

Tell your doctor immediately if you become pregnant or are planning to become pregnant. Your doctor will review your treatment. Do not discontinue Juluca without consulting your doctor, as this may harm you and your unborn child.

### **Breast-feeding**

**Women who are HIV-positive must not breast feed** because HIV infection can be passed on to the baby in breast milk.

**It is not known** whether the ingredients of Juluca can pass into your breast milk.

If you are breast-feeding, or thinking about breast-feeding:

→ **Talk to your doctor immediately.**

### **Driving and using machines**

Juluca can make you dizzy, tired or drowsy and have other side effects that make you less alert.

→ Don't drive or operate machinery unless you are sure you're not affected.

### **Juluca contains lactose**

If you have been told by your doctor that you have intolerance to some sugars, speak with your doctor before taking this medicine.

## **3. How to take Juluca**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- The recommended dose of Juluca is **one tablet once a day**. Juluca **must be taken with a meal**. A meal is important to get the right levels of medicine in your body. A protein-rich nutritional drink alone does not replace a meal.
- Do not chew, crush or split the tablet, to ensure the full dose is taken.

#### Rifabutin

Rifabutin, a medicine to treat some bacterial infections, can lower the amount of Juluca in your body and make it less effective.

If you take rifabutin, your doctor may need to give you an additional dose of rilpivirine. Take the rilpivirine tablet at the same time you take Juluca.

→ Talk to your doctor for further advice on taking rifabutin with Juluca.

#### Antacid medicines

Antacids, to treat indigestion and heartburn, can stop Juluca being absorbed into your body and make it less effective.

Do not take an antacid during the 6 hours before you take Juluca, or for at least 4 hours after you take it.

→ Talk to your doctor for further advice on taking acid-lowering medicines with Juluca.

#### Calcium supplements, iron supplements or multivitamins

Calcium supplements, iron supplements or multivitamins can stop Juluca being absorbed into your body and make it less effective.

Calcium supplements, iron supplements or multivitamins must be taken at the same time as Juluca. Juluca must be taken with a meal.

If you can't take these supplements at the same time as Juluca, do not take calcium supplements, iron supplements or multivitamins during the 6 hours before you take Juluca, or for at least 4 hours after you take it.

→ Talk to your doctor for further advice on taking calcium supplements, iron supplements or multivitamins with Juluca.

#### H<sub>2</sub> receptor antagonists (for example cimetidine, famotidine, nizatidine, ranitidine)

H<sub>2</sub> receptor antagonist medicines can stop Juluca being absorbed into your body and make it less effective. Do not take these medicines during the 12 hours before you take Juluca, or for at least 4 hours after you take it.

→ Talk to your doctor for further advice on taking these medicines with Juluca.

#### **If you take more Juluca than you should**

If you take too many tablets of Juluca **contact your doctor or pharmacist immediately**. If possible, show them the Juluca pack.

#### **If you forget to take Juluca**

If you notice within 12 hours of the time you usually take Juluca, you must take the tablet as soon as possible. The Juluca tablet must be taken with a meal. Then take the next dose as usual. If you notice after 12 hours, then skip that dose and take the next doses as usual.

→**Do not take a double dose** to make up for a forgotten dose.

If you vomit less than 4 hours after taking Juluca, take another tablet with a meal. If you vomit more than 4 hours after taking Juluca you do not need to take another tablet until your next scheduled dose.

#### **Don't stop taking Juluca without advice from your doctor**

Take Juluca for as long as your doctor recommends. Don't stop unless your doctor tells you to.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them, **so it is very important to talk to your doctor about any changes in your health.**

### Allergic reactions

Juluca contains dolutegravir. Dolutegravir can cause a serious allergic reaction known as a *hypersensitivity reaction*. This is an uncommon (may affect up to 1 in 100 people) reaction in people taking dolutegravir. If you get any of the following symptoms:

- skin rash
  - a high temperature (*fever*)
  - lack of energy (*fatigue*)
  - swelling, sometimes of the face or mouth (*angioedema*), causing difficulty in breathing
  - muscle or joint aches
- **See a doctor straight away.** Your doctor may decide to carry out tests to check your liver, kidneys or blood, and may tell you to stop taking Juluca.

### Very common side effects

These may affect **more than 1 in 10 people**:

- headache
- dizziness
- diarrhoea
- feeling sick (*nausea*)
- difficulty in sleeping (*insomnia*).

Very common side effects that may show up in blood tests are:

- increase in the level of liver enzymes (aminotransferases)
- increase in cholesterol
- increase in pancreatic amylase (a digestive enzyme).

### Common side effects

These may affect **up to 1 in 10 people**:

- loss of appetite
- rash
- itching (*pruritus*)
- being sick (*vomiting*)
- stomach (*abdominal*) pain or discomfort
- wind (*flatulence*)
- feeling drowsy
- sleep disorders
- abnormal dreams
- lack of energy (*fatigue*)
- depression (feelings of deep sadness and unworthiness)
- depressed mood
- anxiety
- dry mouth.

Common side effects that may show up in blood tests are:

- increase in the level of enzymes produced in the muscles (creatinine phosphokinase).
- decreased number of platelets, which are involved in blood clotting
- low white blood cell count
- decrease in haemoglobin
- increase in triglycerides (a type of fat)
- increase in lipase (an enzyme involved in breaking down fats)
- increase in bilirubin (a test of liver function) in your blood.

## **Uncommon side effects**

These may affect **up to 1 in 100 people**:

- allergic (*hypersensitivity*) reaction (see ‘allergic reactions’ earlier in this section)
- inflammation of the liver (*hepatitis*)
- suicidal thoughts and behaviours (particularly in patients who have had depression or mental health problems before)
- joint pain
- muscle pain.

## **Rare side effects**

These may affect **up to 1 in 1000 people**:

- liver failure (signs may include yellowing of the skin and the whites of the eyes or unusually dark urine).

## **Not known**

Frequency cannot be estimated from the available data:

- signs or symptoms of inflammation or infection, for example fever, chills, sweats (*immune reactivation syndrome*).

## **Other possible side effects**

People taking combination therapy for HIV may get other side effects.

### **Symptoms of infection and inflammation**

People with advanced HIV infection (AIDS) have weak immune systems, and are more likely to develop serious infections (*opportunistic infections*). Symptoms of infection may develop, caused by old, hidden infections flaring up again as the body fights them. Symptoms usually include **fever**, plus some of the following:

- headache
- stomach ache
- difficulty breathing.

In rare cases, as the immune system becomes stronger, it can also attack healthy body tissue (*autoimmune disorders*). The symptoms of autoimmune disorders may develop many months after you start taking medicine to treat your HIV infection. Symptoms may include:

- palpitations (rapid or irregular heartbeat) or tremor
- hyperactivity (excessive restlessness and movement)
- weakness beginning in the hands and feet and moving up towards the trunk of the body.

**If you get any symptoms of infection** or if you notice any of the symptoms above:

→ **Tell your doctor immediately.** Don’t take other medicines for the infection without your doctor’s advice.

### **Joint pain, stiffness and bone problems**

Some people taking combination therapy for HIV develop a condition called *osteonecrosis*. With this condition, parts of the bone tissue die because of reduced blood supply to the bone. People may be more likely to get this condition:

- if they have been taking combination therapy for a long time
- if they are also taking anti-inflammatory medicines called corticosteroids
- if they drink alcohol
- if their immune systems are very weak
- if they are overweight.

**Signs of osteonecrosis include:**

- stiffness in the joints
- aches and pains in the joints (especially in the hip, knee or shoulder)
- difficulty moving.

If you notice any of these symptoms:

→ **Tell your doctor.**

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Juluca**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and bottle after EXP. The expiry date refers to the last day of that month.

Store in the original package in order to protect from moisture. Keep the bottle tightly closed. Do not remove the desiccant.

This medicine does not require any special temperature storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Juluca contains**

- The active substances are dolutegravir and rilpivirine. Each tablet contains dolutegravir sodium equivalent to 50 mg dolutegravir and rilpivirine hydrochloride equivalent to 25 mg rilpivirine.
- The other ingredients are mannitol (E421), magnesium stearate, microcrystalline cellulose, povidone (K29/32), sodium starch glycolate, sodium stearyl fumarate, lactose monohydrate, croscarmellose sodium, povidone (K30), polysorbate 20, silicified microcrystalline cellulose, polyvinyl alcohol- part hydrolysed, titanium dioxide (E171), macrogol, talc, iron oxide yellow (E172), iron oxide red (E172).
- This medicine contains less than 1 mmol sodium (23 mg) per dosage unit, that is to say essentially 'sodium-free'.

### **What Juluca looks like and contents of the pack**

Juluca film-coated tablets are pink, oval, biconvex tablets debossed with 'SV J3T' on one side.

The film-coated tablets are provided in bottles closed with child-resistant closures.

Each bottle contains 30 film-coated tablets and a desiccant to reduce moisture. Once the bottle has been opened keep the desiccant in the bottle, do not remove it.

Multipacks containing 90 film-coated tablets (3 packs of 30 film-coated tablets) are also available. Not all pack sizes may be available in your country.

### **Marketing Authorisation Holder**

ViiV Healthcare BV  
Huis ter Heideweg 62  
3705 LZ Zeist  
Netherlands

### **Manufacturer**

Glaxo Wellcome, S.A.  
Avda. Extremadura, 3

09400 Aranda De Duero  
Burgos  
Spain

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

**België/Belgique/Belgien**  
ViiV Healthcare srl/bv  
Tél/Tel: + 32 (0) 10 85 65 00

**България**  
ГлаксоСмитКлейн ЕООД  
Тел.: + 359 2 953 10 34

**Česká republika**  
GlaxoSmithKline, s.r.o.  
Tel: + 420 222 001 111  
[cz.info@gsk.com](mailto:cz.info@gsk.com)

**Danmark**  
GlaxoSmithKline Pharma A/S  
Tlf: + 45 36 35 91 00  
[dk-info@gsk.com](mailto:dk-info@gsk.com)

**Deutschland**  
ViiV Healthcare GmbH  
Tel.: + 49 (0)89 203 0038-10  
[viiv.med.info@viivhealthcare.com](mailto:viiv.med.info@viivhealthcare.com)

**Eesti**  
GlaxoSmithKline Eesti OÜ  
Tel: + 372 6676 900  
[estonia@gsk.com](mailto:estonia@gsk.com)

**Ελλάδα**  
GlaxoSmithKline Μονοπρόσωπη Α.Ε.Β.Ε.  
Τηλ: + 30 210 68 82 100

**España**  
Laboratorios ViiV Healthcare, S.L.  
Tel: + 34 900 923 501  
[es-ci@viivhealthcare.com](mailto:es-ci@viivhealthcare.com)

**France**  
ViiV Healthcare SAS  
Tél.: + 33 (0)1 39 17 69 69  
[Infomed@viivhealthcare.com](mailto:Infomed@viivhealthcare.com)

**Hrvatska**  
GlaxoSmithKline d.o.o.  
Tel: + 385 1 6051 999

**Lietuva**  
GlaxoSmithKline Lietuva UAB  
Tel: + 370 5 264 90 00  
[info.lt@gsk.com](mailto:info.lt@gsk.com)  
**Luxembourg/Luxemburg**  
ViiV Healthcare srl/bv  
Belgique/Belgien  
Tél/Tel: + 32 (0) 10 85 65 00

**Magyarország**  
GlaxoSmithKline Kft.  
Tel.: + 36 1 225 5300

**Malta**  
GlaxoSmithKline (Malta) Limited  
Tel: + 356 21 238131

**Nederland**  
ViiV Healthcare BV  
Tel: + 31 (0)30 6986060  
[contact-nl@viivhealthcare.com](mailto:contact-nl@viivhealthcare.com)

**Norge**  
GlaxoSmithKline AS  
Tlf: + 47 22 70 20 00

**Österreich**  
GlaxoSmithKline Pharma GmbH  
Tel: + 43 (0)1 97075 0  
[at.info@gsk.com](mailto:at.info@gsk.com)

**Polska**  
GSK Services Sp. z o.o.  
Tel.: + 48 (0)22 576 9000

**Portugal**  
VIIHVHIV HEALTHCARE, UNIPESSOAL, LDA  
Tel: + 351 21 094 08 01  
[viiv.fi.pt@viivhealthcare.com](mailto:viiv.fi.pt@viivhealthcare.com)

**România**  
GlaxoSmithKline (GSK) S.R.L.  
Tel: + 4021 3028 208

**Ireland**  
GlaxoSmithKline (Ireland) Limited  
Tel: + 353 (0)1 4955000

**Ísland**  
Vistor hf.  
Sími: +354 535 7000

**Italia**  
ViiV Healthcare S.r.l  
Tel: + 39 (0)45 9212611

**Kύπρος**  
GlaxoSmithKline (Cyprus) Ltd  
Τηλ: + 357 22 397000  
gskcyprus@gsk.com

**Latvija**  
GlaxoSmithKline Latvia SIA  
Tel: + 371 67312687  
lv-epasts@gsk.com

**Slovenija**  
GlaxoSmithKline d.o.o.  
Tel: + 386 (0)1 280 25 00  
medical.x.si@gsk.com

**Slovenská republika**  
GlaxoSmithKline Slovakia s. r. o.  
Tel: + 421 (0)2 48 26 11 11  
recepacia.sk@gsk.com

**Suomi/Finland**  
GlaxoSmithKline Oy  
Puh/Tel: + 358 (0)10 30 30 30  
Finland.tuoteinfo@gsk.com

**Sverige**  
GlaxoSmithKline AB  
Tel: + 46 (0)8 638 93 00  
info.produkt@gsk.com

**United Kingdom**  
ViiV Healthcare UK Limited  
Tel: + 44 (0)800 221441  
customercontactuk@gsk.com

**This leaflet was last revised in.**

#### **Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.